

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
23 August 2001 (23.08.2001)

PCT

(10) International Publication Number  
**WO 01/60377 A2**

(51) International Patent Classification<sup>7</sup>: **A61K 31/66**,  
38/43, A61P 3/00, 37/04

Northrop Street, Hopedale, MA 01747 (US). **GOLD-  
BERG, Mark**; 41 Grasmere Road, Needham, MA 02494  
(US).

(21) International Application Number: PCT/US01/03875

(22) International Filing Date: 6 February 2001 (06.02.2001)

(74) Agent: **LAZAR, Steven, R.**; Genzyme Corporation, Legal  
Department, One Kendall Square, Cambridge, MA 02139  
(US).

(25) Filing Language: English

(81) Designated States (*national*): AU, CA, JP.

(26) Publication Language: English

(84) Designated States (*regional*): European patent (AT, BE,  
CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC,  
NL, PT, SE, TR).

(30) Priority Data:  
60/183,296 17 February 2000 (17.02.2000) US  
60/260,069 5 January 2001 (05.01.2001) US

**Published:**

(71) Applicant: **GENZYME CORPORATION** [US/US];  
One Kendall Square, Cambridge, MA 02139 (US).

— without international search report and to be republished  
upon receipt of that report



(72) Inventors: **ZIEGLER, Robin**; 138 Chace Hill Road,  
Sterling, MA 01564 (US). **CHENG, Seng**; 50 Wall Street,  
Wellesley, MA 02184 (US). **MARSHALL, John**; 25

*For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.*

**WO 01/60377 A2**

(54) Title: METHODS FOR TREATMENT OF LYSOSOMAL STORAGE DISEASES

(57) Abstract: Methods are disclosed for treatment of patients suffering from accumulation of a metabolite within macrophages, such as in lysosomal storage diseases. The methods comprise treating the patient with a macrophage depleting amount of a bis-phosphonate compound, such that apoptosis of macrophages is induced and the metabolite is released into circulation so that the metabolites may be eliminated from the patient. The methods may further include administration of a gene therapy vector for the treatment of lysosomal storage diseases.

## METHODS FOR TREATMENT OF LYSOSOMAL STORAGE DISEASES

### FIELD OF THE INVENTION

The present invention relates to methods for improved treatment of lysosomal storage diseases and other medical conditions. The methods include both methods of enzyme replacement therapy and gene therapy.

### BACKGROUND OF THE INVENTION

#### Lysosomal Storage Diseases

Several of the over thirty known lysosomal storage diseases (LSDs) are known to involve a similar pathogenesis, namely, a compromised lysosomal hydrolase. Generally, the activity of a single lysosomal hydrolytic enzyme is reduced or lacking altogether, usually due to inheritance of an autosomal recessive mutation. As a consequence, the substrate of the compromised enzyme accumulates undigested in lysosomes, producing severe disruption of cellular architecture and various disease manifestations.

Gaucher's disease is the oldest and most common lysosomal storage disease known. Type 1 is the most common among three recognized clinical types and follows a chronic course which does not involve the nervous system. Types 2 and 3 both have a CNS component, the former being an acute infantile form with death by age two and the latter a subacute juvenile form. The incidence of Type 1 Gaucher's disease is about one in 50,000 live births generally and about one in 400 live births among Ashkenazim (*see generally* Kolodny et al., 1998, "Storage Diseases of the Reticuloendothelial System", In: Nathan and Oski's Hematology of Infancy and Childhood, 5th ed., vol. 2, David G. Nathan and Stuart H. Orkin, Eds., W.B. Saunders Co., pages 1461-1507). Also known as glucosylceramide lipidosis, Gaucher's disease is caused by inactivation of the enzyme glucocerebrosidase and accumulation of glucocerebroside. Glucocerebrosidase normally catalyzes the hydrolysis of glucocerebroside to glucose and ceramide. In Gaucher's disease, glucocerebroside accumulates in tissue macrophages which become engorged and are typically found in liver, spleen and bone marrow and occasionally in lung, kidney and intestine. Secondary hematologic sequelae include severe anemia and thrombocytopenia in addition to the characteristic progressive hepatosplenomegaly and skeletal complications, including osteonecrosis and osteopenia with secondary pathological fractures.

Niemann-Pick disease, also known as sphingomyelin lipidosis, comprises a group of disorders characterized by foam cell infiltration of the reticuloendothelial system. Foam cells in

Niemann-Pick become engorged with sphingomyelin and, to a lesser extent, other membrane lipids including cholesterol. Niemann-Pick is caused by inactivation of the enzyme sphingomyelinase in Types A and B disease, with 27-fold more residual enzyme activity in Type B (see Kolodny et al., 1998, *Id.*). The pathophysiology of major organ systems in Niemann-Pick can be briefly summarized as follows. The spleen is the most extensively involved organ of Type A and B patients. The lungs are involved to a variable extent, and lung pathology in Type B patients is the major cause of mortality due to chronic bronchopneumonia. Liver involvement is variable, but severely affected patients may have life-threatening cirrhosis, portal hypertension, and ascites. The involvement of the lymph nodes is variable depending on the severity of disease. Central nervous system (CNS) involvement differentiates the major types of Niemann-Pick. While most Type B patients do not experience CNS involvement, it is characteristic in Type A patients. The kidneys are only moderately involved in Niemann Pick disease.

Fabry disease is an X-linked recessive LSD characterized by a deficiency of  $\alpha$ -galactosidase A ( $\alpha$ -Gal A), also known as ceramide trihexosidase, which leads to vascular and other disease manifestations via accumulation of glycosphingolipids with terminal  $\alpha$ -galactosyl residues, such as globotriaosylceramide (GL-3) (see generally Desnick RJ et al., 1995,  $\alpha$ -Galactosidase A Deficiency: Fabry Disease, In: The Metabolic and Molecular Bases of Inherited Disease, Scriver et al., eds., McGraw-Hill, New York, 7<sup>th</sup> ed., pages 2741-2784). Symptoms may include anhidrosis (absence of sweating), painful fingers, left ventricular hypertrophy, renal manifestations, and ischemic strokes. The severity of symptoms varies dramatically (Grewal RP, 1994, Stroke in Fabry's Disease, J. Neurol. 241, 153-156). A variant with manifestations limited to the heart is recognized, and its incidence may be more prevalent than once believed (Nakao S, 1995, An Atypical Variant of Fabry's Disease in Men with Left Ventricular Hypertrophy, N. Engl. J. Med. 333, 288-293).

Recognition of unusual variants can be delayed until quite late in life, although diagnosis in childhood is possible with clinical vigilance (Ko YH et al., 1996, Atypical Fabry's Disease - An Oligosymptomatic Variant, Arch. Pathol. Lab. Med. 120, 86-89; Mendez MF et al., 1997, The Vascular Dementia of Fabry's Disease, Dement. Geriatr. Cogn. Disord. 8, 252-257; Shelley ED et al., 1995, Painful Fingers, Heat Intolerance, and Telangiectases of the Ear: Easily Ignored Childhood Signs of Fabry Disease, Pediatric Derm. 12, 215-219). The mean age of diagnosis of Fabry disease is 29 years.

Replacement of the defective enzyme is considered feasible using a recombinant retrovirus carrying the cDNA encoding  $\alpha$ -Gal A to transfect skin fibroblasts obtained from Fabry patients (Medin JA et al., 1996, Correction in Trans for Fabry Disease: Expression, Secretion, and Uptake of  $\alpha$ -Galactosidase A in Patient-Derived Cells Driven by a High-Titer Recombinant Retroviral Vector, Proc. Natl. Acad. Sci. USA 93, 7917-7922).

### SUMMARY OF THE INVENTION

Accordingly, the present invention provides methods for the treatment of lysosomal storage diseases and other conditions. The methods may comprise therapy to deplete macrophages or Kupffer cells as a therapy or adjunct therapy for eliminating lysosomal storage products which might otherwise be stored in macrophages and/or Kupffer cells. In addition or as an alternative, macrophage depletion or apoptosis of macrophages or Kupffer cells may be performed prior to or contemporaneously with enzyme replacement therapy and/or gene therapy which utilize DNA molecules which encode a therapeutic protein of interest, such as glucocerebrosidase or sphingomyelinase, under conditions suitable for the expression of said DNA molecule.

Thus, in certain embodiments, the present invention comprises methods of treating a patient suffering from accumulation of a metabolite within macrophages, said method comprising treating the patient with a macrophage depleting amount of a bisphosphonate compound, such that apoptosis of macrophages is induced and the metabolite is released into circulation so that it may be eliminated from the patient. In preferred embodiments, the patient may be suffering from Gaucher's disease, in which case the metabolite is GL1; or the patient may be suffering from Niemann-Pick disease, in which case the metabolite is sphingomyelin. In further preferred embodiments, the methods further comprise administering to the patient a composition of purified recombinant protein. For patients with Gaucher's disease, the recombinant protein is recombinant glucocerebrosidase. For patients with Niemann-Pick disease, the recombinant protein is recombinant acid sphingomyelinase.

In other embodiments, the present invention comprises methods of treating a patient suffering from accumulation of a metabolite within macrophages, said method comprising treating the patient with a macrophage depleting amount of a bisphosphonate compound, such that apoptosis of macrophages is induced, and administering to the patient a gene therapy vector encoding a compound which is able to break down the accumulated metabolite.

In preferred embodiments, the patient may be suffering from Gaucher's disease, and the gene therapy vector encodes glucocerebrosidase. Optionally, the method may further comprise administering to the patient a composition of purified recombinant glucocerebrosidase. In embodiments wherein the patient is suffering from Niemann-Pick disease, the gene therapy  
5 vector encodes acid sphingomyelinase, and the method may optionally further comprise administering to the patient a composition of purified recombinant acid sphingomyelinase.

In still other embodiments of the present invention, the patient is suffering from Fabry's disease, and the gene therapy vector encodes alpha galactosidase A. Optionally, the method may further comprise administering to the patient a composition of purified recombinant alpha-  
10 galactosidase. In yet other embodiments, the patient is suffering from Pompe disease, and the gene therapy vector encodes alpha glucosidase. Optionally, the method may further comprise administering to the patient a composition of purified recombinant alpha glucosidase. In another embodiment, the patient is suffering from Hurler's Disease (MPS-I), and the gene therapy vector encodes alpha-L iduronidase. Optionally, the method may further comprise administering to the  
15 patient a composition of purified recombinant alpha-L iduronidase.

In the most preferred embodiments, the macrophage depleting compound is a bisphosphonate, such as clodronate, alendronate, pamidronate, zoledronate, etidronate, ibandronate, olpadronate, risedronate, medronate, neridronate, tiludronate, or incadronate. Other apoptosis inducing agents may be used, appropriately targeted to macrophages, such as through  
20 encapsulation in liposomes. In addition, the methods of the present invention may employ other compounds which will exhibit a macrophage-depleting or macrophage-inhibiting effect. By "macrophage-depleting" or "macrophage-inhibiting" effect, it is meant that the compound is present in sufficient amount to alter or block the activity of these cells and to thereby reduce or eliminate endocytosis or phagocytosis. Such compounds or substances may include, for  
25 example, doxorubicin, gamma globulin, heavy metal salts and molecules which may block the Fc receptor.

The present methods have important advantages for the treatment of lysosomal storage diseases. First, the methods of the present invention allow the persistent expression of therapeutic levels of lysosomal storage enzymes to be produced from gene therapy vectors.  
30 Second, the methods of the present invention allow for the elimination of significant amounts of lysosomal storage products which might otherwise be sequestered within macrophages and/or Kupffer cells. Third, the methods of the present invention may allow for more effective

treatment of lysosomal storage diseases using enzyme replacement therapy and/or gene therapy in which lower dosage regimens may conveniently be used.

The methods of the present invention are particularly adapted towards the treatment of lysosomal storage diseases in which the lysosomal storage product is typically concentrated within macrophages. It is believed that by depleting the macrophages, the accumulated lysosomal storage product will enter the system and to a significant extent will be eliminated from the system through a combination of secretion in bile and excretion in urine, rather than stored in the macrophages. These include Gaucher's disease and Niemann-Pick Disease. Although the associated lysosomal enzymes are not as strongly localized to macrophages in other lysosomal storage diseases, such treatment of such diseases may be enhanced by the methods of the present invention. These include LAL, Pompe's (alpha-glucosidase), Hurler's (MPS I) (alpha-L iduronidase), Fabry's (alpha-galactosidase), Hunters (MPS II) (iduronate sulfatase), Morquio Syndrome (MPS IVA) (galactosamine-6-sulfatase), MPS IVB, (beta-galactosidase) and Maroteux-Lamy C (MPS VI)(arylsulfatase B). In addition, any disease in which an unwanted metabolite is accumulated within macrophages or Kupffer cells, or any therapy for which depletion of macrophages might otherwise be desired, may be a suitable candidate for therapy using the methods of the present invention.

The preferred coding DNA sequences useful for gene therapy targeting to the liver via depletion of macrophages include DNA sequences which encode a therapeutic protein for which activity and targeting to liver is desired. By depletion of macrophages, and particularly Kupffer cells, it is believed that more of the gene therapy vector can be taken up by hepatocytes, where they can efficiently produce higher levels of therapeutic protein. In particular, preferred coding DNA sequences include those sequences encoding, glucocerebrosidase and acid sphingomyelinase, for the treatment of patients with Gaucher's Disease and Niemann-Pick Disease, respectively. Other preferred coding DNA sequences include those encoding alpha-glucosidase (Pompe's Disease), alpha-L iduronidase (Hurler's Disease or MPS I), alpha-galactosidase (Fabry Disease), iduronate sulfatase (Hunters Disease (MPS II), galactosamine-6-sulfatase (MPS IVA), beta galactosidase (MPS IVB) and arylsulfatase B (MPS VI).

As described further herein, the methods of the present invention, which utilize macrophage-depleting agents such as clodronate, the macrophage-depleting agent may be utilized in combination with other agents which have a macrophage-depleting or macrophage-inhibiting effect. By macrophage-inhibiting effect, it is intended to mean that the substance is

able to interact with macrophages, either via the Fc receptor or other means, and is thereby able to reduce or eliminate the activity of such macrophages in endocytosis or phagocytosis.

In either aspect of the present invention, treatment with clodronate or other macrophage-depleting substance, may be useful in conjunction with more traditional therapies, such as enzyme-replacement therapy. Thus, for the treatment of Gaucher disease, the methods of the present invention may be used in addition to treatment with recombinantly produced glucocerebrosidase, commercially available as Cerezyme® [Genzyme Corporation, Cambridge, MA; also see United States Patent 5,236,838]. For treatment of Fabry disease, the methods of the present invention may be used in addition to treatment with recombinantly produced alpha-galactosidase [see United States Patent 5,580,757]. Use of the methods of the present invention may allow for the use of lower doses, or less frequent dosing, with enzyme replacement therapy.

#### BRIEF DESCRIPTION OF THE FIGURES

**Figure 1:** Enhanced expression is generated from a low dose of virus by pre-treating mice with clodronate liposomes to deplete Kupffer cells in the liver.

Groups of four mice were given the following three treatment regimes: A)  $1 \times 10^{11}$  particles of Ad2/CMVHI $\alpha$ gal, B)  $2 \times 10^{10}$  particles of Ad2/CMVHI $\alpha$ gal, C)  $2 \times 10^{10}$  particles of Ad2/CMVHI $\alpha$ gal mixed with  $8 \times 10^{10}$  particles Ad2/EV. Two additional groups were pre-treated 24 hours prior to administration of a low dose of virus, one with 50  $\mu$ l clodronate liposomes to deplete macrophages in the liver (Kupffer cells), the other with 50  $\mu$ l PBS liposomes as a control. Mice were sacrificed three days following virus administration. Liver homogenates were analyzed for  $\alpha$ -galactosidase A expression using an ELISA specific for human  $\alpha$ -galactosidase A. Expression from a low dose of Ad2/CMVHI $\alpha$ gal was enhanced in mice pre-treated with clodronate liposomes. The effect was similar to that seen in the group supplemented with empty vector.

**Figure 2:** Dose-dependant expression is achieved following macrophage depletion by clodronate liposomes.

Groups of BALB/c mice were injected with various doses of Ad2/CMVHI $\alpha$ gal as described in the graph. 24 hours prior to virus administration one half of the mice at each dose of the vector had been treated with 50  $\mu$ l of clodronate liposomes to deplete macrophages in the liver (Kupffer cells). Tissue homogenates were analyzed for  $\alpha$ -galactosidase A expression using the ELISA (the shaded area with the graph represents the range of  $\alpha$ -galactosidase A in normal mouse



tissues). Three tissues were evaluated for expression as shown. Mice treated with clodronate liposomes showed higher levels of expression at all doses compared to mice receiving virus alone. The expression levels were closely correlated with the dose of virus administered. Therapeutic levels of  $\alpha$ -galactosidase A were attained with a 50-fold lower dose of virus after clodronate liposome treatment.

**Figure 3: Immunohistochemical staining of liver sections from mice treated with clodronate liposomes.**

Mice from the experiment described in Figure 2 were perfused with 10% neutral-buffered formalin upon sacrifice, the livers removed and fixed for 24 hours. The samples were embedded in paraffin and 0.5  $\mu$ m sections cut. Sections were deparaffinized, rehydrated and immunohistochemistry performed. Sections were stained with a rat monoclonal antibody to the mouse macrophage surface marker F4/80 (Serotec) using the VectaStain ABC kit from Vector Laboratories. Slides were counterstained from hematoxylin and visualized at 200x magnification. Macrophages can be identified by the dark brown stain. Kupffer cells were almost completely depleted in the livers of mice treated with clodronate liposomes.

**Figure 4: Improved persistence and efficacy from a reduced dose of adenovirus following macrophage depletion**

$1 \times 10^9$  particles of virus were injected into the tail vein of two groups of Fabry mice. One group had been pre-treated with clodronate liposomes. Organs were divided to be assayed for both  $\alpha$ -galactosidase A expression and GL-3 levels. Tissues were assayed by ELISA. Figure 4A shows  $\alpha$ -galactosidase A levels. The shaded area within the graph represents the range of  $\alpha$ -galactosidase A in normal mouse tissues. Figure 4B shows GL-3 levels. The dashed lines represent mice pre-treated with clodronate liposomes, and the solid lines represent mice treated with virus alone. Values represent an average of four treated mice per group. Clodronate liposome pre-treatment enhanced levels and persistence of expression from  $1 \times 10^9$  particles of Ad2/CMVHI $\alpha$ gal with resulting GL-3 clearance in all tissues except kidney. This dose of vector was not sufficient to clear GL-3 in Fabry mice treated with virus alone.

**Figure 5: Antibody levels in plasma from Fabry mice treated with Ad2/CMVHI $\alpha$ gal +/- clodronate liposomes**

Antibody titers were measured in the plasma samples collected from the animals described in Figure 4. Antibodies made to human  $\alpha$ -galactosidase were assayed using an ELISA format. Symbols represent titers from individual animals. **Fabry mice pre-treated with clodronate liposomes made significantly lower levels of antibodies to  $\alpha$ -galactosidase A than mice treated with adenovirus alone.**

#### **DETAILED DESCRIPTION OF THE INVENTION**

It is known that many of the lysosomal enzymes, and in particular, glucocerebrosidase (Gaucher's Disease) and acid sphingomyelinase (Niemann-Pick Disease A & B), are particularly good candidates for methods of treatment involving depletion of macrophages, since the lysosomal storage products involved in these diseases in large part accumulate in the macrophages and Kupffer cells.

Other lysosomal storage disorders which may be suitable for gene therapy avoidance of macrophages and Kupffer cells include LAL, Pompe's (alpha-glucosidase), Hurler's (alpha-L iduronidase), Fabry's (alpha-galactosidase), Hunters (MPS II) (iduronate sulfatase), Morquio Syndrome (MPS IVA)(galactosamine-6-sulfatase), MPS IVB (beta-D-galactosidase), and Maroteux-Lamy C (MPS VI)(arylsulfatase B).

There is evidence of other independent pathways, in addition to the mannose-6-phosphate pathway, that may function in the transport of lysosomal enzymes inside cells and of alternate mechanisms for the internalization of lysosomal enzymes by cell-surface receptors in addition to mannose-6-phosphate receptors (Scriver et al. 1995). In addition, any protein for which avoidance of the macrophages is desired may be a suitable candidate for the method of the present invention.

In certain embodiments of the present invention, purified recombinant proteins are produced in cell culture, using coding DNA sequences to transduce cells, such as CHO cells, yeast cells, or other cells suitable for recombinant protein production. In other embodiments of the present invention, nucleotide sequences encoding a lysosomal enzyme or other therapeutic protein are provided via administration of gene therapy vectors. Such gene therapy vectors include recombinant viral vectors containing coding DNA sequences, in which aspects of viruses are used to transfect cells, which then will express the coding DNA sequences to produce the encoded lysosomal enzyme or therapeutic protein. Suitable virus systems are known for example, for adenovirus, adeno-associated virus and other parvoviruses, alphaviruses, and

retroviruses, including lentiviruses. Other gene therapy vectors include non-viral systems for transducing cells, such as the use of cationic lipids or other cationic molecules as the cell-transduction mediator. Also known in the art are methods for delivery of naked DNA into cells.

The preferred coding DNA sequences include sequences encoding any therapeutic protein. In preferred embodiments, the coding DNA sequence is one which is desired to be targeted to hepatocytes, without accumulation in macrophages. In particular, preferred coding DNA sequences include those sequences encoding, glucocerebrosidase for the treatment of patients with Gaucher's Disease and acid sphingomyelinase for the treatment of patients with Niemann-Pick Disease, respectively. Other preferred coding DNA sequences include those encoding alpha-glucosidase (Pompe's Disease), alpha-L iduronidase (Hurler's Disease), alpha-galactosidase (Fabry's Disease), and iduronate sulfatase (Hunters Disease (MPS II), galactosamine-6-sulfatase (MPS IVA); beta-D-galactosidase (MPS IVB); and arylsulfatase B (MPS VI). In addition, a cDNA for any protein to be expressed from liver, for which avoidance of the macrophages is desired may be a suitable candidate for the methods of the present invention.

Methods for the purification of recombinant human proteins are well-known, including methods for the production of recombinant human glucocerebrosidase [for Gaucher's Disease]; sphingomyelinase [for Niemann-Pick Disease], alpha-galactosidase [for Fabry Disease]; alpha-glucosidase [for Pompe's Disease]; alpha-L iduronidase [for Hurler's Syndrome]; iduronate sulfatase [for Hunter's Syndrome]; galactosamine-6-sulfatase [for MPS IVA]; beta-D-galactosidase [for MPS IVB]; and arylsulfatase B [for MPS VI]. See, for example, Scriver et al., eds., The Metabolic and Molecular Bases of Inherited Diseases, Vol. II., 7<sup>th</sup> ed. (McGraw-Hill, NY; 1995), the disclosure of which is hereby incorporated herein by reference.

Compounds which may be useful in the methods of the present invention include compounds which are able to induce apoptosis of macrophages and Kupffer Cells. Such compounds include members of the bisphosphonate family, such as clodronate, alendronate, pamidronate, zoledronate, etidronate, ibandronate, olpadronate, risedronate, medronate, neridronate, tiludronate, and incadronate. [See Rogers et al., Cancer, 88/12 Suppl. 2961-2978 (2000)]. Other apoptosis-inducing agents which may be useful in the methods of the present invention include Fas ligand [see Watanabe-Fukunaga et al., *Nature* 314:317, 1992; Zhou et al., *J. Exp. Med.* 176:1063-1072, 1992; Nagata, *Adv. Immunol.* 57:129-144, 1994, and Dhein et al., *Nature* 373:438-441, 1995]. In general, a wide range of apoptosis inducing agent concentrations

can be used. Van Rooijen and Sanders, (1994) J. Immunological Methods 174:83-93, identifies preferred concentrations for clodronate encapsulated in liposomes. See also, Van Rooijen and Sanders (1996) Hepatology 23:1239-1243; Wang et al (1999) Oral Pathol. Med. 28:145-151; Schugart et al. (1999) Gene Therapy, 6:448-453; Lieber et al. (1997), J. Virology, 71:8798-8807; 5 Kuzmin et al. (1997) Gene Therapy, 4:309-316; Worgall et al. (1997), Human Gene Therapy, 8:1675-1684; Stein et al. (1998), Gene Therapy, 5:431-439; Wolff et al. (1997) J. Virology, 71:624-629; Van Rooijen et al. (1996) J. Immunological Methods, 193:93-99; and Van Rooijen and Sanders (1997) Trends in Biotechnology, 15:178-185.

#### 10 Treatment of Gaucher's Disease and Niemann-Pick Disease – Apoptosis of Kupffer Cells

Therapy for metabolic storage diseases of macrophages [e.g., Gaucher disease] may be accomplished by the use of agents that disrupt macrophages, such as bisphosphonates. In Gaucher disease, the primary clinical manifestation is massive accumulation of glucosyl ceramide, or GL1, the substrate for the enzyme glucocerebrosidase [GCR], in the macrophages, 15 particularly of the liver or the spleen. Induction of apoptosis in these macrophages, by treatment with bisphosphonates such as clodronate or compounds with similar effects would cause release of the accumulated glycolipid. This could allow for removal of GL1 from the body through bile salt secretion and/or urine excretion. Circulating GL1 may also re-distribute into cell types which are more readily accessible to treatment by enzyme replacement therapy and/or gene 20 therapy either by direct administration into and/or transduction of the affected cells, or by the affected cells' ability to take up circulating mature GCR secreted by a distantly transduced site [e.g., a targeted cell or depot organ].

Circulating GL1 must be primarily reabsorbed by hepatocytes in order for it to be targeted for biliary secretion. Indications are that the majority of a bolus injection of liposome- 25 encapsulated GL1 ends up in the liver parenchyma. Pentchev et al., BBA, (1981); Tokoro et al., J. Lip. Res., (1987); Hers, Gastroenterology 48:625-632 (1965). An additional amount is found in Kupffer Cells. It appears that rapid influx of circulating GL1 storage product, and the loss of liver Kupffer Cells are not harmful to patients. As a result, it is predicted by the inventors that GL1 can be eliminated from the body of Gaucher's Disease patients by apoptosis of the 30 Kupffer Cells, potentially in combination with enzyme replacement treatment and/or gene therapy treatment with GCR.



transduced depot organ.

Treatment of other lysosomal storage disorders, such as Fabry Disease, may require significant transduction of a target depot organ, such as liver or lung. Secretion of the therapeutic transgene is driven by overexpression. Secretion into circulation and subsequent uptake by distal diseased tissues is required for efficacy of this treatment. We have shown in Fabry mice that a high dose of adenoviral vector ( $10^{11}$  particles) was required for efficient clearance of the lipidic substrate deposited in diseased tissues, globotriaosylceramide, or GL-3. This dose of adenovirus can be associated with significant liver toxicity. Reducing the dose of adenovirus particles resulted in a non-linear dose response in expression; a ten-fold reduction in dose yielded a 150 fold drop in expression. Treating mice systemically with clodronate liposomes depleted macrophages from the liver and boosted the level of expression, thus therapeutic levels of expression were attained from a 50-fold lower dose of virus. We propose that this effect can extend to other therapeutic proteins for lysosomal storage diseases and other diseases; to other depot organs, such as lung; as well as to other gene therapy vectors, such as adeno-associated virus, other viral-based gene therapy vectors, and non-viral gene therapy vectors. Clodronate liposome treatment, or any other method for circumventing macrophage uptake of vectors, should allow us to treat effectively with a lower, less toxic dose of gene therapy vector.

30 Fabry mice are pre-treated with liposomes 24 hours prior to gene therapy vector administration to deplete macrophages. 'Liposomes' may comprise:

- a. Clodronate liposomes prepared as described . Van Rooijen and Sanders, (1994) J. Immunological Methods 174:83-93.
- b. Liposomes encapsulating other drugs effective at depleting macrophages without activation of said macrophages.

5        In addition, gene therapy vectors encoding a lysosomal enzyme, such as alpha-galactosidase in the case of Fabry Disease, may be complexed with molecules (such as PEG) which further shield them from recognition and uptake by macrophages.

**Alternative or Additional Macrophage-Depletion Strategies:**

10        In addition to, or instead of, treatment with clodronate or other bisphosphonates, other methods are available which may prove useful for the depletion of macrophages and/or Kupffer cells, and thus for reducing or avoiding the immune response to gene therapy, in accordance with the methods of the present invention. These include use of other macrophage-depleting  
15        substances, such as doxorubicin encapsulated in liposomes, which has been demonstrated to have a macrophage depleting effect. Daemen et al, Int. J. Cancer. 61:716-721 (1995). Doxorubicin, which is approved for cancer therapy, has some associated toxicities. However, by using lower and less frequent doses, it may be possible to achieve a macrophage depleting effect which aids in prolonged expression from gene therapy vectors, without severe adverse effects. Other  
20        molecules which might be useful for methods of the present invention include molecules which may specifically induce apoptosis or ablation of macrophages or Kupffer cells. This may include Fas ligand or other apoptotic or ablative agents targeted to macrophages. One preferred method for targeting macrophages is encapsulation in liposomes.

25        As described further herein, the methods of the present invention, which utilize macrophage-depleting agents such as clodronate, the macrophage-depleting agent may be utilized in combination with other agents which have a macrophage-depleting or macrophage-inhibiting effect. By macrophage-inhibiting effect, it is intended to mean that the substance is able to interact with macrophages, either via the Fc receptor or other means, and is thereby able to reduce or eliminate the activity of such macrophages in endocytosis or phagocytosis.

30        In either aspect of the present invention, treatment with clodronate or other macrophage-depleting substance, may be useful in conjunction with more traditional therapies, such as enzyme-replacement therapy. Thus, for the treatment of Gaucher disease, the methods of the present invention may be used in addition to treatment with recombinantly produced

glucocerebrosidase, commercially available as Cerezyme® [Genzyme Corporation, Cambridge, MA; also see United States Patent 5,236,838]. For treatment of Fabry disease, the methods of the present invention may be used in addition to treatment with recombinantly produced alpha-galactosidase [see United States Patent 5,580,757]. Use of the methods of the present invention  
5 may allow for the use of lower doses, or less frequent dosing, with enzyme replacement therapy.

As an alternative, or additional, strategy to effectively deplete macrophages or Kupffer cells, approaches can be used which will alter or block the activity of these cells to thereby reduce or eliminate endocytosis or phagocytosis. A number of drugs are available in this regard.  
10 Most notably, human gamma globulins (IG) may be used. This class of drugs is presently approved product for supplementing immune system. While the exact mechanism of action is not determined, it is believed that these drugs can essentially swamp the immune system by forming a non-specific reticuloendothelial system blockade. See Samuelsson et al., Science 291:484-486 (2001).

15 Methods of the invention may further comprise the use of other methods include the use of molecules which may be able to block or mask the Fc receptors on macrophages. Such molecules may include heavy metal salts, such as gadolinium chloride [see Husztik et al., *Brit. J. Exper. Path.*, 61:624-30 (1980); and Hardonk et al., *J. Leukocyte Biol.*, 52:296-302 (1992)], derivatives of deoxynojirimycin [DNJ] [see Overkleef et al., *J. Biol. Chem.*, 273:26522-26527  
20 (1998); WO98/02161] a sugar analog inhibitor of glucosylceramide synthase; and mannin, which is essentially a polymer of numerous mannose entities. Other such substances may include large neutral polymers such as polyethylene glycol (PEG) or hyaluronic acid (HA) polymers.

\* \* \* \*

While the invention is exemplified with respect to the production of specific proteins,  
25 these examples are not to be interpreted as limiting the invention in any manner. The invention is further not limited to any stated belief as to the mechanism for action. As described above, and as will be clear to those skilled in the art from reading the specification, the methods of the present invention are useful for the treatment of lysosomal storage diseases, and for a number of other conditions which may involve accumulation of unwanted substances within macrophages,  
30 including the lysosomal storage products described above. Many modifications and variations of the methods and materials used in the present description will also be apparent to those skilled in the art. Such modifications and variations fall within the scope of the invention.



The entire disclosures of all of the publications and references cited in this specification are hereby incorporated herein by reference.

### EXAMPLES

5

#### **Viral-mediated gene therapy of Fabry disease.**

Fabry disease is caused by a deficiency of the lysosomal hydrolase  $\alpha$ -galactosidase A. This results in progressive accumulation of globotriaosylceramide (GL-3) in lysosomes. The feasibility of using gene therapy for treating Fabry disease has been demonstrated. Intravenous  
10 administration of Ad2/CMVHI $\alpha$ gal into Fabry mice resulted in correction of both the enzymatic and lysosomal storage defects in all affected organs. However there was associated liver toxicity and expression subsided over several months. The methods of the present invention will be useful in combination with more advanced versions of adenoviral vectors with improved toxicity profiles, as well as strategies to enhance delivery of the virus to hepatocytes. In order to  
15 moderate toxicity, we sought to lower the dose of vector administered. However, reducing the dose of Ad2/CMVHI $\alpha$ gal by ten fold resulted in a greater than 200 fold drop in expression of  $\alpha$ -galactosidase A, a level that was insufficient to completely clear accumulated GL-3. In order to address this problem, mice were pre-treated with clodronate liposomes to deplete Kupffer Cells. Administration of  $2 \times 10^{10}$  particles of Ad2/CMVHI $\alpha$ gal in the presence of clodronate liposomes  
20 generated higher expression levels of  $\alpha$ -galactosidase A than in control animals that were treated with PBS liposomes. The level of  $\alpha$ -galactosidase A attained in the presence of clodronate was consistent with the dose used. This suggests that sequestration of viral particles by the Kupffer cells may play a role in the non-linear dose response. The use of clodronate liposomes resulted in lower levels of antibodies forming to human  $\alpha$ -galactosidase A. Using the lower doses of  
25 adenoviral particles, which are effective persistent expression with administration of clodronate-containing liposome, results in reduced levels of cytokines, such as IL-6 and IL-12, which are associated with toxicity of high doses of adenoviral particles. Hence, strategies such as the use of clodronate, for the depletion of Kupffer cells to prevent this process, may allow the use of lower therapeutic doses of gene therapy vectors for the treatment of lysosomal storage diseases,  
30 such as Fabry, and ultimately safer gene therapy treatment.

An alternative viral vector reported to show lower toxicity and more persistent expression is adeno-associated virus (AAV). Systemic administration of AAV/CMVHI $\alpha$ gal into immune-

suppressed Fabry mice generated  $\alpha$ -galactosidase A levels in the liver that were approximately 10% of normal. These levels were undiminished over three months. Although expression was significantly lower than the levels obtained from adenovirus vectors, GL-3 was reduced in all tissues assayed except kidney. Thus, depletion of macrophages using the methods of the present invention may also be applicable to AAV vectors. In addition, AAV vectors containing optimized transcription cassettes may be used in order to enhance expression to levels that are therapeutic for use in Fabry disease.

#### Viral-mediated gene therapy of Gaucher disease.

We have used clodronate in two studies using adenoviral vectors encoding the human glucocerebrosidase (GC) gene. The experimental details are essentially the same as for  $\alpha$ gal except we used wild-type BALB/c mice as there presently is no viable *in vivo* Gaucher disease model.

In the first study, we analyzed the levels of GC enzyme activity obtained from a tail vein intravenous bolus delivery of  $10^{11}$  virus particles in 100 $\mu$ l. There were 3 groups of 4 mice (BALB/c) each. Group A: naïve mice. Group B: received virus, without clodronate. Group C received a dose of clodronate the day before virus administration. Resulting expression compared to naïve was:

Group A. serum 1x; liver 1x

Group B. serum 12x (33 $\mu$ g/ml); liver 3x (96 $\mu$ g/gm)

Group C. serum 500x (1400 $\mu$ g/ml); liver 230x (750 $\mu$ g/gm)

The second experiment was a repeat of the first but included two additional groups, each group receiving a viral dose of  $2 \times 10^{10}$ , Group D without clodronate; Group E with clodronate.

Resulting expression compared to naïve was:

Group A. serum 1x; liver 1x

Group B. serum 1x; liver 9x (131 $\mu$ g/gm)

Group C. serum 1600x (7520 $\mu$ g/ml); liver 117x (1209 $\mu$ g/gm)

Group D. serum 1x; liver 4x (58 $\mu$ g/gm)

Group E. serum 117x (553 $\mu$ g/ml); liver 72x (1053 $\mu$ g/gm)

In summary, mice that received  $10^{11}$  Ad2/hGC particles exhibited liver expression of GC protein at ~5x endogenous levels, and serum GC levels at ~6x endogenous levels. Pre-treatment with clodronate liposomes yielded liver expression on the order of 100-fold that of endogenous GC levels and serum GC levels on the order of 1000-fold that of endogenous GC levels


5 following the same dose of virus.

We claim:

1. A method of treating a patient suffering from accumulation of a metabolite within macrophages, said method comprising treating the patient with a macrophage depleting amount of a bisphosphonate compound, such that apoptosis of macrophages is induced and  
5 the metabolite is released into circulation so that it may be eliminated from the patient.
2. The method of claim 1, wherein the bisphosphonate compound is clodronate.
3. The method of claim 1, wherein the patient is suffering from Gaucher's disease, and the metabolite is GL1.
4. The method of claim 3, further comprising administering to the patient a composition of  
10 purified recombinant glucocerebrosidase.
5. The method of claim 1, wherein the patient is suffering from Niemann-Pick disease, and the metabolite is sphingomyelin.
6. The method of claim 5, further comprising administering to the patient a composition of purified recombinant acid sphingomyelinase.
- 15 7. A method of treating a patient suffering from accumulation of a metabolite within macrophages, said method comprising treating the patient with a macrophage depleting amount of a bisphosphonate compound, such that apoptosis of macrophages is induced, and administering to the patient a gene therapy vector encoding a compound which is able to break down the accumulated metabolite.
- 20 8. The method of claim 7, wherein the patient is suffering from Gaucher's disease, and the gene therapy vector encodes glucocerebrosidase.
9. The method of claim 8, further comprising administering to the patient a composition of purified recombinant glucocerebrosidase.
10. The method of claim 7, wherein the patient is suffering from Niemann-Pick disease, and the  
25 the gene therapy vector encodes acid sphingomyelinase

11. The method of claim 10, further comprising administering to the patient a composition of purified recombinant acid sphingomyelinase.
12. The method of claim 7, wherein the patient is suffering from Fabry's disease, and the gene therapy vector encodes alpha galactosidase A.
- 5 13. The method of claim 12, further comprising administering to the patient a composition of purified recombinant alpha-galactosidase.
14. The method of claim 7, wherein the patient is suffering from Pompe disease, and the gene therapy vector encodes alpha glucosidase.
15. The method of claim 14, further comprising administering to the patient a composition of  
10 purified recombinant alpha glucosidase.
16. The method of claim 7, wherein the patient is suffering from Hurler's Disease (MPS-I), and the gene therapy vector encodes alpha-L iduronidase.
17. The method of claim 16, further comprising administering to the patient a composition of purified recombinant alpha-L iduronidase.
- 15 18. The method of claim 7, further comprising administration of a macrophage depleting or macrophage inhibiting compound selected from the group consisting of doxycirubin, gamma globulin, and neutral polymers.

1/6



IV LIPOSOMES 24 hr PRE-VIRUS

FIG. 1

2/6


DOSE-DEPENDENT EXPRESSION  
FOLLOWING MACROPHAGE DEPLETION BY CLODRONATE LIPOSOMES

FIG. 2A




FIG. 2B




FIG. 2C

3/6




FIG. 3B




FIG. 3A



$\alpha$ -GALACTOSIDASE A LEVELS FOLLOWING LOW DOSE ADENOVIRUS  
ADMINISTRATION IN CLODRONATE TREATED FABRY MICE




FIG. 4A

5/6





FIG. 4B

6/6



\*n = 3 WITH 1 VERY HIGH OUTLIER

FIG. 5

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
23 August 2001 (23.08.2001)

PCT

(10) International Publication Number  
**WO 01/60377 A3**

- (51) International Patent Classification<sup>7</sup>: A61K 31/66, 38/43, 48/00, A61P 3/00, 37/04 Northrop Street, Hopedale, MA 01747 (US). GOLD-  
BERG, Mark; 41 Grasmere Road, Needham, MA 02494 (US).
- (21) International Application Number: PCT/US01/03875 (74) Agent: LAZAR, Steven, R.; Genzyme Corporation, Legal  
Department, One Kendall Square, Cambridge, MA 02139 (US).
- (22) International Filing Date: 6 February 2001 (06.02.2001)
- (25) Filing Language: English (81) Designated States (*national*): AU, CA, JP.
- (26) Publication Language: English (84) Designated States (*regional*): European patent (AT, BE,  
CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC,  
NL, PT, SE, TR).
- (30) Priority Data:  
60/183,296 17 February 2000 (17.02.2000) US  
60/260,069 5 January 2001 (05.01.2001) US  
Published:  
— with international search report
- (71) Applicant: GENZYME CORPORATION [US/US]; One Kendall Square, Cambridge, MA 02139 (US). (88) Date of publication of the international search report:  
18 April 2002
- (72) Inventors: ZIEGLER, Robin; 138 Chace Hill Road, Sterling, MA 01564 (US). CHENG, Seng; 50 Wall Street, Wellesley, MA 02184 (US). MARSHALL, John; 25
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*



WO 01/60377 A3

(54) Title: METHODS FOR TREATMENT OF LYSOSOMAL STORAGE DISEASES USING BIPHOSPHONATES

(57) Abstract: Methods are disclosed for treatment of patients suffering from accumulation of a metabolite within macrophages, such as in lysosomal storage diseases. The methods comprise treating the patient with a macrophage depleting amount of a bisphosphonate compound, such that apoptosis of macrophages is induced and the metabolite is released into circulation so that the metabolites may be eliminated from the patient. The methods may further include administration of a gene therapy vector for the treatment of lysosomal storage diseases.

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 01/03875

<b>A. CLASSIFICATION OF SUBJECT MATTER</b> IPC 7 A61K31/66 A61K38/43 A61K48/00 A61P3/00 A61P37/04		
According to International Patent Classification (IPC) or to both national classification and IPC		
<b>B. FIELDS SEARCHED</b> Minimum documentation searched (classification system followed by classification symbols) IPC 7 A61K A61P		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used) EPO-Internal, MEDLINE, WPI Data, PAJ, BIOSIS, CHEM ABS Data, EMBASE		
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	VAN ROOIJEN N. ET AL.: "Transient suppression of macrophage functions by liposome-encapsulated drugs." TRENDS IN BIOTECHNOLOGY., vol. 15, 1997, pages 178-185, XP002175114 * see page 180 left col., page 181 right col. and pages 182-183 *	1-18
Y	STEIN C. S. ET AL.: "Systemic and central nervous system correction of lysosomal storage in mucopolysaccharidosis type VII mice." J. VIROL., vol. 73, no. 4, 1999, pages 3424-3429, XP002175115 * see abstract, page 3424 right col. and page 3427 *	1-18
<div style="display: flex; justify-content: space-between;"> <span><input checked="" type="checkbox"/> Further documents are listed in the continuation of box C.</span> <span><input checked="" type="checkbox"/> Patent family members are listed in annex.</span> </div>		
<div style="display: flex;"> <div style="flex: 1;"> <p>* Special categories of cited documents:</p> <p>*A* document defining the general state of the art which is not considered to be of particular relevance</p> <p>*E* earlier document but published on or after the international filing date</p> <p>*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>*O* document referring to an oral disclosure, use, exhibition or other means</p> <p>*P* document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="flex: 1;"> <p>*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>*G* document member of the same patent family</p> </div> </div>		
Date of the actual completion of the international search  16 August 2001		Date of mailing of the international search report  07/09/2001
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl. Fax: (+31-70) 340-3016		Authorized officer  Merckling, V

## INTERNATIONAL SEARCH REPORT

International Application No.

PCT/US 01/03875

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 98 02161 A (VIANELLO PAOLA ;KOOMEN GERRIT JAN (NL); UNIV AMSTERDAM (NL); AERTS) 22 January 1998 (1998-01-22) * see claim 16, pages 11-12 and 23 *	1-18
Y	ZIEGLER R J ET AL: "CORRECTION OF ENZYMATIC AND LYSOSOMAL STORAGE DEFECTS IN FABRY MICE BY ADENOVIRUS-MEDIATED GENE TRANSFER" HUMAN GENE THERAPY, XX, XX, vol. 10, no. 10, 1 July 1999 (1999-07-01), pages 1667-1682, XP001012918 ISSN: 1043-0342 * see abstract and page 1679 *	1-18
A	"PARTICULARITIES OF LYSOSOMAL STORAGE DISEASES GENE THERAPY" GENE THERAPY, MACMILLAN PRESS LTD., BASINGSTOKE, GB, vol. 3, no. 12, December 1996 (1996-12), pages 1039-1041, XP001007386 ISSN: 0969-7128 * see pages 1040-1041 *	1-18
Y	STEIN C S ET AL: "EFFECTS OF MACROPHAGE DEPLETION AND ANTI-CD40 LIGAND ON TRANSGENE EXPRESSION AND REDOSING WITH RECOMBINANT ADENOVIRUS" GENE THERAPY, MACMILLAN PRESS LTD., BASINGSTOKE, GB, vol. 5, no. 4, April 1998 (1998-04), pages 431-439, XP001008505 ISSN: 0969-7128 * see abstract and page 435 *	1-18
X	BEMBI B ET AL: "Aminohydroxypropylidene- biphosphonate in the treatment of bone lesions in a case of Gaucher 's disease type 3." ACTA PAEDIATRICA, (1994 JAN) 83 (1) 122-4. XP001008466	1-4
Y	* see abstract *	1-18
A	"The Merck Manual, 17th edition." 1999, MERCK RESEARCH LABORATORIES XP002175116 * pages 214-215, 2388 and 2408 *	1-18

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 01/03875

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9802161 A	22-01-1998	AU 732552 B	26-04-2001
		AU 3464797 A	09-02-1998
		EP 0912179 A	06-05-1999
		JP 2000516577 T	12-12-2000
		US 6177447 B	23-01-2001
<hr/>			